

Positive Solutions of Three-Point Boundary Value Problems for the One-Dimensional p -Laplacian with Infinitely Many Singularities

BING LIU

Department of Mathematics
Huazhong University of Science and Technology
Wuhan 430074, Hubei, P.R. China
bingliug@public.wh.hb.cn

(Received September 2002; revised and accepted September 2003)

Abstract—We consider the singular three-point boundary value problems

$$\begin{aligned}(\phi_p(y'))' + a(t)f(y(t)) &= 0, & 0 < t < 1, \\ y'(0) = 0, \quad y(1) &= \beta y(\eta),\end{aligned}$$

where $\phi_p(s) = |s|^{p-2}s$, $p \geq 2$, $0 < \beta < 1$, $0 < \eta < 1$, $f \in C([0, +\infty), [0, +\infty))$, $a : [0, 1] \rightarrow [0, +\infty)$, and has countably many singularities in $[0, 1/2)$. We show that there exist countably many positive solutions by using the fixed-point index theory. © 2004 Elsevier Ltd. All rights reserved.

Keywords—Singularity, One-dimensional p -Laplacian, Multiple positive solutions, Three-point boundary value problems, Fixed-point index.

1. INTRODUCTION

We are interested in the existence of positive solutions of the following singular three-point boundary value problems for one-dimensional p -Laplacian:

$$(\phi_p(y'))' + a(t)f(y(t)) = 0, \quad 0 < t < 1, \quad (1.1)$$

$$y'(0) = 0, \quad y(1) = \beta y(\eta), \quad (1.2)$$

where $\phi_p(s) = |s|^{p-2}s$, $p \geq 2$, $0 < \beta < 1$, $0 < \eta < 1$, $f \in C([0, +\infty), [0, +\infty))$, $a : [0, 1] \rightarrow [0, +\infty)$, and has countably many singularities in $[0, 1/2)$.

Recently, for the existence problems of positive solutions of multipoint boundary value problems for second-order ordinary differential equations, some authors have obtained the existence results. For details, see, for example, [1–5]. However, the multipoint boundary value problems treated in the above-mentioned references do not discuss problems with singularities. For the singular case of multipoint boundary value problems, with the author's knowledge, no one has studied the

This project is supported by Postdoctoral Science Foundation of China (No. 200114).

existence of positive solutions in the case. Very recently, Kaufmann and Kosmatov [6] showed that there exist countably many positive solutions for the two-point boundary value problems with infinitely many singularities of following form:

$$u''(t) + b(t)f(u(t)) = 0, \quad 0 < t < 1, \quad (1.3)$$

$$u(0) = u(1) = 0, \quad (1.4)$$

where $b \in L^p[0, 1]$ for some $p \geq 1$ and has countably many singularities in $[0, 1/2)$.

Motivated by the result of [6], we show that problems (1.1), (1.2) have infinitely many solutions if a and f satisfy some suitable conditions. The key tool in our approach is the following fixed-point index theorem [7].

THEOREM A. (See [7].) *Let E be Banach a space and $P \subset E$ be a cone in E . Let $r > 0$ define $\Omega_r = \{x \in P \mid \|x\| < r\}$. Assume that $A : \bar{\Omega} \rightarrow P$ is a completely continuous operator such that $Ax \neq x$ for $x \in \partial\Omega_r$.*

(i) *If $\|Au\| \leq \|u\|$ for $u \in \partial\Omega_r$, then*

$$i(A, \Omega_r, P) = 1.$$

(ii) *If $\|Au\| \geq \|u\|$ for $u \in \partial\Omega_r$, then*

$$i(A, \Omega_r, P) = 0.$$

From now on, in the rest of the paper, we always assume that $a(t)$ satisfies the following conditions.

(H₀) There exists a sequence $\{t_i\}_{i=1}^\infty$ such that $t_{i+1} < t_i$ ($i \in N$), $t_1 < 1/2$, $\lim_{i \rightarrow \infty} t_i = t^* \geq 0$, $\lim_{t \rightarrow t_i} a(t) = +\infty$ for all $i = 1, 2, \dots$, and

$$0 < \int_0^1 a(s) ds < +\infty.$$

Moreover, $a(t)$ does not vanish identically on any subinterval of $[0, 1]$.

Note, it is easy to check that Condition (H₀) implies that

$$0 < \int_0^1 \phi_q \left(\int_0^s a(s_1) ds_1 \right) ds < +\infty,$$

where $\phi_q(s)$ is the inverse function to $\phi_p(s)$, a.e., $\phi_q(s) = |s|^{q-2}s$, $1/p + 1/q = 1$.

2. MAIN RESULT AND REMARK

Let $E = C[0, 1]$, and only the supnorm is used. Denote

$$C_0^+[0, 1] = \left\{ y \in E : \min_{t \in [0, 1]} y(t) \geq 0, y'(0) = 0, y(1) = \beta y(\eta) \right\},$$

$$P = \{ y \in C_0^+[0, 1] : y(t) \text{ is a nonnegative concave function} \}.$$

It is obvious that P is a cone in E .

LEMMA 2.1. Suppose (H_0) holds. Then, the function

$$A(t) = \frac{\beta}{1-\beta} \int_t^{1-t_1} \phi_q \left(\int_t^s a(s_1) ds_1 \right) ds + \int_{t_1}^t \phi_q \left(\int_{t_1}^s a(s_1) ds_1 \right) ds, \quad t \in [t_1, 1-t_1],$$

is continuous and positive on $[t_1, 1-t_1]$. Furthermore, $L(t_1) = \min_{t \in [t_1, 1-t_1]} A(t) > 0$.

PROOF. At first, it is easily seen that $A(t)$ is continuous on $[t_1, 1-t_1]$. Next, let

$$A_1(t) = \frac{\beta}{1-\beta} \int_t^{1-t_1} \phi_q \left(\int_t^s a(s_1) ds_1 \right) ds \quad \text{and} \quad A_2(t) = \int_{t_1}^t \phi_q \left(\int_{t_1}^s a(s_1) ds_1 \right) ds.$$

Then, from Condition (H_0) , we know that the function $A_1(t)$ is strictly monotone decreasing on $[t_1, 1-t_1]$ and $A_1(1-t_1) = 0$, the function $A_2(t)$ is strictly monotone increasing on $[t_1, 1-t_1]$ and $A_2(t_1) = 0$. So the function $A(t) = A_1(t) + A_2(t_1)$ is positive on $[t_1, 1-t_1]$, which implies $L(t_1) = \min_{t \in [t_1, 1-t_1]} A(t) > 0$.

LEMMA 2.2. Let $y \in P$ and $\mu \in (0, 1/2)$. Then,

$$y(t) \geq \mu \|y\|, \quad t \in [\mu, 1-\mu],$$

where $\|y\| = \sup_{t \in [0,1]} y(t)$.

PROOF. Let

$$\tau = \inf \left\{ \xi \in [0, 1] : \sup_{t \in [0,1]} y(t) = y(\xi) \right\}.$$

CASE (i). $\tau \in [0, \mu]$. It follows from the concavity of $y(t)$ that each point on chord between $(\tau, y(\tau))$ and $(1, y(1))$ is below the graph of $y(t)$. Thus,

$$y(t) \geq y(\tau) + \frac{y(1) - y(\tau)}{1 - \tau} (t - \tau), \quad t \in [\mu, 1 - \mu].$$

Hence,

$$\begin{aligned} y(t) &\geq \min_{t \in [\mu, 1-\mu]} \left[y(\tau) + \frac{y(1) - y(\tau)}{1 - \tau} (t - \tau) \right] \\ &= y(\tau) + \frac{y(1) - y(\tau)}{1 - \tau} (1 - \mu - \tau) \\ &= \frac{1 - \mu - \tau}{1 - \tau} y(1) + \frac{\mu}{1 - \tau} y(\tau) \geq \mu y(\tau), \end{aligned}$$

which implies that

$$y(t) \geq \mu \|y\|.$$

CASE (ii). $\tau \in [\mu, 1-\mu]$. If $t \in [\mu, \tau]$, similarly, we have

$$y(t) \geq y(\tau) + \frac{y(\tau) - y(0)}{\tau} (t - \tau), \quad t \in [\mu, \tau].$$

Thus,

$$\begin{aligned} y(t) &\geq \min_{t \in [\mu, \tau]} \left[y(\tau) + \frac{y(\tau) - y(0)}{\tau} (t - \tau) \right] \\ &= y(\tau) + \frac{y(\tau) - y(0)}{\tau} (\mu - \tau) \\ &= \frac{\mu}{\tau} y(\tau) + \left(1 - \frac{\mu}{\tau} \right) y(0) \geq \mu y(\tau). \end{aligned}$$

If $t \in [\tau, 1 - \mu]$, similarly,

$$y(t) \geq y(\tau) + \frac{y(1) - y(\tau)}{1 - \tau}(t - \tau), \quad t \in [\tau, 1 - \mu].$$

Thus,

$$\begin{aligned} y(t) &\geq \min_{t \in [\tau, 1 - \mu]} \left[y(\tau) + \frac{y(1) - y(\tau)}{1 - \tau}(t - \tau) \right] \\ &= y(\tau) + \frac{y(1) - y(\tau)}{1 - \tau}(1 - \mu - \tau) \\ &= \frac{\mu}{1 - \tau}y(\tau) + \frac{1 - \mu - \tau}{1 - \tau}y(1) \geq \mu y(\tau). \end{aligned}$$

Therefore, we obtain

$$y(t) \geq \mu \|y\|, \quad t \in [\mu, 1 - \mu].$$

CASE (iii). $\tau \in [1 - \mu, 1]$. Similarly, we have

$$y(t) \geq y(\tau) + \frac{y(\tau) - y(0)}{\tau}(t - \tau), \quad t \in [\mu, 1 - \mu].$$

Thus,

$$\begin{aligned} y(t) &\geq \min_{t \in [\mu, 1 - \mu]} \left[y(\tau) + \frac{y(\tau) - y(0)}{\tau}(t - \tau) \right] \\ &= y(\tau) + \frac{y(\tau) - y(0)}{\tau}(\mu - \tau) \\ &= \frac{\mu}{\tau}y(\tau) + \left(1 - \frac{\mu}{\tau}\right)y(0) \geq \mu y(\tau), \end{aligned}$$

which yields

$$y(t) \geq \mu \|y\|, \quad t \in [\mu, 1 - \mu].$$

This completes the proof.

Now we define an operator $T : P \rightarrow P$ by

$$(Ty)(t) = \frac{\beta}{1 - \beta} \int_{\eta}^1 \phi_q \left(\int_0^s a(s_1) f(x(s_1)) ds_1 \right) ds + \int_t^1 \phi_q \left(\int_0^s a(s_1) f(x(s_1)) ds_1 \right) ds. \quad (2.1)$$

Then, it is easy to see that $(Ty)(t) \geq 0$ ($0 \leq t \leq 1$), $(Ty)'(0) = 0$, $(Ty)(1) = \beta(Ty)(\eta)$, and $(Ty)'(t) = -\phi_q(\int_0^t a(s) f(x(s)) ds) \leq 0$, $(\phi_p(Ty)'(t))' = -a(t)f(y(t)) \leq 0$. This shows that $T(P) \subset P$.

LEMMA 2.3. Let (H_0) hold, then $T : P \rightarrow P$ is completely continuous.

PROOF. Let D be a bounded subset of P and $M > 0$ is the constant such that $\|y\| \leq M$ for $y \in D$. Then, we

$$\|Ty\| \leq \left[\frac{1}{1 - \beta} \int_0^1 \phi_q \left(\int_0^s a(s_1) ds_1 \right) ds \right] \phi_q(\sup\{f(y) : y \in D\}),$$

which implies the boundedness of $T(D)$.

Now, for any $y \in D$, $0 \leq t_1 < t_2 \leq 1$, we have

$$\begin{aligned} |(Ty)(t_1) - (Ty)(t_2)| &= \left| \int_{t_1}^{t_2} \phi_q \left(\int_0^s a(s_1) f(x(s_1)) ds_1 \right) ds \right| \\ &\leq \phi_q \left(\int_0^1 a(s) ds \right) \phi_q(\sup\{f(y) : y \in D\}) |t_1 - t_2|. \end{aligned}$$

Therefore, $T(D)$ is equicontinuous.

Finally, in view of the continuity of f and the Lebesgue's dominated convergence theorem, it is easy to see that T is continuous on D . Thus, the Arzela-Ascoli theorem implies that $T : P \rightarrow P$ is completely continuous.

THEOREM 2.1. Assume that Condition (H_0) holds. Let $\{\mu_k\}_{k=1}^\infty$ such that $\mu_k \in (t_{k+1}, t_k)$ ($k = 1, 2, \dots$). Let $\{R_k\}_{k=1}^\infty$ and $\{r_k\}_{k=1}^\infty$ such that

$$R_{k+1} < \mu_k r_k < r_k < \Lambda_1 r_k < R_k, \quad k = 1, 2, \dots,$$

where $\Lambda_1 \in (1/L(t_1), +\infty)$.

Furthermore, for each natural number k , assume that f satisfies

$$(H_1) \quad f(u) \geq (\Lambda_1 r_k)^{p-1} \text{ for all } u \in [\mu_k r_k, r_k];$$

$$(H_2) \quad f(u) \leq (\Lambda_2 R_k)^{p-1} \text{ for all } u \in [0, R_k], \text{ where } 0 < \Lambda_2 < (((1-\beta\eta)/(1-\beta))\phi_q(\int_0^1 a(s) ds))^{-1}.$$

Then, the problems (1.1), (1.2) have infinitely many positive solutions $\{u_i\}_{i=1}^\infty$ such that

$$r_i \leq \|u_i\| \leq R_i, \quad \text{for each } i = 1, 2, \dots$$

PROOF. Since $t^* < t_{k+1} < \mu_k < t_k < 1/2$ for all $k \in N$. Then, for each $k \in N$ and $y \in P$, from Lemma 2.2, we have

$$y(t) \geq \mu_k \|y\|, \quad t \in [\mu_k, 1 - \mu_k]. \quad (2.2)$$

Consider the sequence $\{\Omega_k^1\}_{k=1}^\infty$ and $\{\Omega_k^2\}_{k=1}^\infty$ of open subsets of $C_0^+[0, 1]$ defined by

$$\Omega_k^1 = \{y \in P : \|y\| < r_k\},$$

$$\Omega_k^2 = \{y \in P : \|y\| < R_k\}.$$

Now, fix k and let $y \in \partial\Omega_k^1$. From (2.2), we have $r_k = \|y\| \geq y(s) \geq \mu_k \|y\| = \mu_k r_k$, for $s \in [\mu_k, 1 - \mu_k]$. In the following, we consider three cases (note that $[t_1, 1 - t_1] \subset [\mu_k, 1 - \mu_k]$).

CASE 1. If $\eta \in [t_1, 1 - t_1]$. In this case, from (2.1), Condition (H_1) , and Lemma 2.1, we get

$$\begin{aligned} \|Ty\| &= (Ty)(0) \\ &= \frac{\beta}{1-\beta} \int_\eta^1 \phi_q \left(\int_0^s a(s_1) f(y(s_1)) ds_1 \right) ds + \int_0^1 \phi_q \left(\int_0^s a(s_1) f(y(s_1)) ds_1 \right) ds \\ &\geq \frac{\beta}{1-\beta} \int_\eta^{1-t_1} \phi_q \left(\int_\eta^s a(s_1) f(y(s_1)) ds_1 \right) ds + \int_{t_1}^\eta \phi_q \left(\int_{t_1}^s a(s_1) f(y(s_1)) ds_1 \right) ds \\ &\geq \Lambda_1 r_k \left[\frac{\beta}{1-\beta} \int_\eta^{1-t_1} \phi_q \left(\int_\eta^s a(s_1) ds_1 \right) ds + \int_{t_1}^\eta \phi_q \left(\int_{t_1}^s a(s_1) ds_1 \right) ds \right] \\ &\geq \Lambda_1 r_k A(\eta) \\ &\geq \Lambda_1 r_k L(t_1) > r_k = \|y\|. \end{aligned}$$

CASE 2. If $\eta \in (0, t_1)$, from (2.1), Condition (H_1) , and Lemma 2.1, we get

$$\begin{aligned} \|Ty\| &= (Ty)(0) \\ &= \frac{\beta}{1-\beta} \int_\eta^1 \phi_q \left(\int_0^s a(s_1) f(y(s_1)) ds_1 \right) ds + \int_0^1 \phi_q \left(\int_0^s a(s_1) f(y(s_1)) ds_1 \right) ds \\ &\geq \frac{\beta}{1-\beta} \int_{t_1}^{1-t_1} \phi_q \left(\int_{t_1}^s a(s_1) f(y(s_1)) ds_1 \right) ds \\ &\geq \Lambda_1 r_k A(t_1) \\ &\geq \Lambda_1 r_k L(t_1) > r_k = \|y\|. \end{aligned}$$

CASE 3. If $\eta \in (1 - t_1, 1)$, from (2.1), Condition (H_1) , and Lemma 2.1, we get

$$\begin{aligned} \|Ty\| &= (Ty)(0) \\ &= \frac{\beta}{1-\beta} \int_\eta^1 \phi_q \left(\int_0^s a(s_1) f(y(s_1)) ds_1 \right) ds + \int_0^1 \phi_q \left(\int_0^s a(s_1) f(y(s_1)) ds_1 \right) ds \\ &\geq \int_{t_1}^{1-t_1} \phi_q \left(\int_{t_1}^s a(s_1) f(y(s_1)) ds_1 \right) ds \\ &\geq \Lambda_1 r_k A(1 - t_1) \\ &\geq \Lambda_1 r_k L(t_1) > r_k = \|y\|. \end{aligned}$$

Thus, in all cases, an application of Theorem A shows that

$$i(T, \Omega_k^1, P_k) = 0. \quad (2.3)$$

Next, let $y \in \partial\Omega_k^2$. Then, from (2.1), Condition (H_2) , we have

$$\begin{aligned} \|Ty\| &= (Ty)(0) \\ &= \frac{\beta}{1-\beta} \int_{\eta}^1 \phi_q \left(\int_0^s a(s_1) f(y(s_1)) ds_1 \right) ds + \int_0^1 \phi_q \left(\int_0^s a(s_1) f(y(s_1)) ds_1 \right) ds \\ &\leq \Lambda_2 R_k \left[\frac{\beta}{1-\beta} \int_{\eta}^1 \phi_q \left(\int_0^s a(s_1) ds_1 \right) ds + \int_0^1 \phi_q \left(\int_0^s a(s_1) ds_1 \right) ds \right] \\ &\leq \Lambda_2 R_k \cdot \frac{1-\beta\eta}{1-\beta} \phi_q \left(\int_0^1 a(s) ds \right) \\ &< R_k = \|y\|. \end{aligned}$$

Thus, Theorem A implies

$$i(T, \Omega_K^2, P_k) = 1. \quad (2.4)$$

Hence, since $r_k < R_k$ for $k \in N$, (2.3) and (2.4), it follows from additivity of the fixed-point index that

$$i(T, \Omega_k^2 \setminus \bar{\Omega}_k^1, P) = 1, \quad \text{for } k \in N.$$

Thus, for each $k \in N$, T has a fixed point in $\Omega_k^2 \setminus \bar{\Omega}_k^1$ such that $r_k \leq \|y\| \leq R_k$. Since $k \in N$ was arbitrary, the proof is complete.

REMARK. There exist many functions $a(t)$ that satisfy Condition (H_0) . For example, let $\Delta = \sqrt{2}(\pi^2/3 - 9/4)$, and define

$$t_0 = \frac{5}{16}, \quad t_n = t_0 - \sum_{i=0}^{n-1} \frac{1}{(i+2)^4}, \quad n = 1, 2, \dots$$

We consider the function $a(t) : [0, 1] \rightarrow (0, +\infty)$ given by

$$a(t) = \sum_{n=1}^{\infty} a_n(t), \quad t \in [0, 1],$$

where

$$a_n(t) = \begin{cases} \frac{2}{n(n+1)(t_{n+1} + t_n)}, & 0 \leq t < \frac{t_{n+1} + t_n}{2}, \\ \frac{1}{\Delta(t_n - t)^{1/2}}, & \frac{t_{n+1} + t_n}{2} \leq t < t_n, \\ \frac{1}{\Delta(t - t_n)^{1/2}}, & t_n < t \leq \frac{t_n + t_{n-1}}{2}, \\ \frac{2}{n(n+1)(2 - t_n - t_{n-1})}, & \frac{t_n + t_{n-1}}{2} < t \leq 1. \end{cases}$$

At first, it is easily seen that $t_1 = 1/4 < 1/2$, $t_n - t_{n+1} = 1/((n+2)^4)$, $n = 1, 2, \dots$, and (note that $\sum_{n=1}^{\infty} 1/n^4 = \pi^4/90$)

$$t^* = \lim_{n \rightarrow \infty} t_n = \frac{5}{16} - \sum_{i=0}^{\infty} \frac{1}{(i+2)^4} = \frac{5}{16} - \left(\frac{\pi^4}{90} - 1 \right) = \frac{21}{16} - \frac{\pi^4}{90} > \frac{1}{5}.$$

Next, since $\sum_{n=1}^{\infty} 1/n^2 = \pi^2/6$, we have

$$\begin{aligned}
 \sum_{n=1}^{\infty} \int_0^1 a_n(t) dt &= \sum_{n=1}^{\infty} \frac{2}{n(n+1)} + \frac{1}{\Delta} \sum_{n=1}^{\infty} \left[\int_{(t_{n+1}+t_n)/2}^{t_n} \frac{1}{(t_n-t)^{1/2}} dt \right. \\
 &\quad \left. + \int_{t_n}^{(t_n+t_{n-1})/2} \frac{1}{(t-t_n)^{1/2}} dt \right] \\
 &= 2 + \frac{\sqrt{2}}{\Delta} \sum_{n=1}^{\infty} \left[(t_n - t_{n+1})^{1/2} + (t_{n-1} - t_n)^{1/2} \right] \\
 &= 2 + \frac{\sqrt{2}}{\Delta} \sum_{n=1}^{\infty} \left[\frac{1}{(n+2)^2} + \frac{1}{(n+1)^2} \right] \\
 &= 2 + \frac{\sqrt{2}}{\Delta} \left[\left(\frac{\pi^2}{6} - \frac{5}{4} \right) + \left(\frac{\pi^2}{6} - 1 \right) \right] \\
 &= 2 + \frac{\sqrt{2}}{\Delta} \left[\frac{\pi^2}{3} - \frac{9}{4} \right] \\
 &= 3.
 \end{aligned}$$

Hence,

$$\int_0^1 a(t) dt = \int_0^1 \sum_{n=1}^{\infty} a_n(t) dt = \sum_{n=1}^{\infty} \int_0^1 a_n(t) dt < \infty,$$

which implies that Condition (H_0) holds.

REFERENCES

1. R. Ma, Positive solutions for a nonlinear three-point boundary value problem, *Electron. J. Diff. Eqns.* **34**, 1–8, (1999).
2. R. Ma and N. Castaneda, Existence of solutions for nonlinear m -point boundary value problems, *J. Math. Anal. Appl.* **256**, 556–567, (2001).
3. J.R.L. Webb, Positive solutions for some three-point boundary value problems via fixed point index, *Nonlinear Anal.* **47**, 4319–4332, (2001).
4. B. Liu, Positive solutions for a nonlinear three-point boundary value problem, *Computers Math. Applic.* **44** (1/2), 201–211, (2002).
5. B. Liu, Positive solutions for a nonlinear three-point boundary value problem, *Appl. Math. Comput.* **132**, 11–28, (2002).
6. E.R. Kaufmann and N. Kosmatov, A multiplicity result for a boundary value problem with infinitely many singularities, *J. Math. Anal. Appl.* **269**, 444–453, (2002).
7. K. Deimling, *Nonlinear Functional Analysis*, Springer, New York, (1985).